

Multi-taxa inventory of naturalized species in Chile

Nicol Fuentes¹, Alicia Marticorena¹, Alfredo Saldaña¹, Viviane Jerez²,
Juan Carlos Ortiz², Pedro Victoriano², Rodrigo A. Moreno^{3,4}, Juan Larraín⁵,
Cristobal Villaseñor-Parada^{6,7}, Götz Palfner¹, Paulina Sánchez^{6,8}, Aníbal Pauchard^{6,8}

1 Departamento de Botánica, Facultad de Ciencias Naturales y Oceanográficas, Universidad de Concepción, Chile **2** Departamento de Zoología, Facultad de Ciencias Naturales y Oceanográficas, Universidad de Concepción, Chile **3** Facultad de Ciencias, Universidad Santo Tomás, Av. Ejército 146, Santiago, Chile **4** Centro de Investigación e Innovación para el Cambio Climático (CIICC), Universidad Santo Tomás, Av. Ejército 146, Santiago, Chile **5** Instituto de Biología, Pontificia Universidad Católica de Valparaíso, Campus Curauma, Av. Universidad 330, Valparaíso, Chile **6** Laboratorio de Invasiones Biológicas (LIB), Facultad de Ciencias Forestales, Universidad de Concepción, Chile **7** Laboratorio de Estudios Algales (ALGALAB), Departamento de Oceanografía, Facultad de Ciencias Naturales y Oceanográficas, Universidad de Concepción, Chile **8** Instituto de Ecología y Biodiversidad (IEB), Santiago, Chile

Corresponding author: Nicol Fuentes (nfuentes@udec.cl)

Academic editor: Marcel Rejmanek | Received 12 June 2020 | Accepted 27 July 2020 | Published 12 August 2020

Citation: Fuentes N, Marticorena A, Saldaña A, Jerez V, Ortiz JC, Victoriano P, Moreno RA, Larraín J, Villaseñor-Parada C, Palfner G, Sánchez P, Pauchard A (2020) Multi-taxa inventory of naturalized species in Chile. *NeoBiota* 60: 25–41. <https://doi.org/10.3897/neobiota.60.55366>

Abstract

Here we present a multi-taxa inventory of naturalized alien species recorded on continental Chile and adjacent marine habitats, including eight taxonomic groups. We identified 1,122 species. These comprise 790 vascular plants (terrestrial and aquatic); 31 nonvascular plants [Bryophyta (mosses), Marchantiophyta (liverworts) and Anthocerotophyta (hornworts)]; 18 marine and freshwater macro and micro algae; 71 fungi; 39 terrestrial vertebrates (amphibians, reptiles, mammals and birds); 108 insects; 37 marine and freshwater invertebrates and vertebrates (6 polychaetes, 3 mollusks and 28 Pisces); and 28 terrestrial gastropods. For all taxonomic groups, naturalized species were found to mainly be distributed in regions with Mediterranean and temperate climates, with few at either extreme of the country. The invasion curves show that naturalized species first underwent a positive increment, followed by an apparent plateau phase,

mainly in vascular plants, insects and vertebrates. In fungi, marine and freshwater macro and microalgae, vertebrates and invertebrates, the cumulative number of naturalized species increased sharply starting in the early 20th century; the lack of collections before 1900 is also evident. When considering naturalized species as a whole, this inventory highlights that the rate of new naturalizations consistently increased after 1950, especially for some taxonomic groups such as insects, fungi, and vascular plants. This multi-taxa inventory of naturalized species provides a platform for national reporting on biodiversity indicators and highlights areas where Chile must invest resources to manage biological invasions.

Keywords

alien species, Chile, database, invasion periods, rate of introduction

Introduction

Inventories of naturalized alien species are not only fundamental to elucidate the causes and consequences of the invasion phenomenon (Mack et al. 2000; Pimentel et al. 2005; van Kleunen et al. 2015; Seebens et al. 2017), but also because of their relevance in nature conservation, ecosystem functioning, human health and economy (Hulme 2009; Pyšek and Richardson 2010). Currently, there is an urgent need for inventories of naturalized species, since they reflect these species' local patterns and provide critical information about invasive species distribution (van Kleunen et al. 2015; Seebens et al. 2017). However, in developing countries, naturalized species distribution is still poorly documented and understood (Gardener et al. 2012; Speziale et al. 2012; but see Castilla et al. 2005; Moreno et al. 2006; Castilla and Neill 2009 for marine non-native species; Fuentes et al. 2013 for plants). These species have been neglected in collections and studies due to a historical bias that found no scientific value in studying the naturalized component (Pauchard et al. 2004; Fuentes et al. 2013). Thus, a national Chilean naturalized species inventory is now essential for the early detection and potential eradication or control of invasive species through risk assessment protocols (Moreno et al. 2006; Fuentes et al. 2010, 2013). Additionally, naturalized species inventories provide datasets suitable for the analysis of temporal patterns of biological invasions, in contrast with the current pattern that reflects geographical biases in information on invasion patterns (Pyšek et al. 2008; Núñez and Pauchard 2010).

Approaches such as invasion curves and variations in the rate of naturalized species introductions allow us to identify invasion periods as well as the temporal dynamics of species accumulations (Fuentes et al. 2008; Seebens et al. 2017). Pyšek and Prach (1993) prepared invasion curves methods for reconstructing the propagation history of four alien species in the Czech Republic. They adjusted an exponential model to the accumulated number of locations against time. The slope of the corresponding regression line was used as a measure of the invasion rate (Mihulka and Pyšek 2001). Abrupt inflexions on the invasion curve indicate expansion periods

of the alien species involved (Pyšek and Prach 1993). In this context, whether different taxa show similar invasion curves in a given region is an open question, hence, comparing invasion curves can reveal differences in the invasion process. Seebens et al. (2017) proposed that differences in the pathways and distribution of species introductions suggest that the chronology of invasion could vary among taxonomic groups. This approach allows us to infer whether the increase in numbers of naturalized species shows any sign of saturation, or whether we can expect biological invasions to continue increasing (Seebens et al. 2017).

A substantial part of Chile has been recognized as a hotspot of world biodiversity (Ormazabal 1993; Myers et al. 2000; Mittermeier et al. 2005) due to its remarkably high levels of endemism and the biogeographic isolation of several taxonomic groups (Armesto et al. 1998; Habit et al. 2006; Vidal and Díaz-Páez 2012; Rodríguez et al. 2018), raising concerns regarding its susceptibility to invasions (Arroyo et al. 2000; Fuentes et al. 2015; Seebens et al. 2017). The extraordinary biogeographic characteristics of Chile make it ideal for understanding biological invasion patterns and have great potential for inferring future invasion trends. However, the study of biological invasions in Chile has been addressed mainly in the last decade, both in terrestrial (Quiroz et al. 2009), and marine systems (Castilla and Neill 2009; Villaseñor-Parada et al. 2017). Therefore, the few inventories that exist of naturalized species have been created for only certain taxonomic groups (e.g., boring polychaetes, Moreno et al. 2006; vascular plants, Fuentes et al. 2013; ascidians, Turon et al. 2016; aquatic plants, Urrutia et al. 2017; marine seaweed, Villaseñor-Parada et al. 2018), with no comprehensive inventory of naturalized species, which would allow for the identification of invasion patterns at the multi-taxa level. To address this knowledge gap, the Project GEF/MMA/PNUD, aimed to develop the first national inventory of naturalized species, including eight taxonomic groups recorded on continental Chile and adjacent marine habitats (PNUD 2017). In this paper, we have updated these lists with current taxonomic status and the full dataset has been included as supplementary material. Here, we present a comprehensive inventory of naturalized species in Chile and analyze the distributional and temporal trends of biological invasions in the country in order to identify priority responses to the growing threat from biological invasions.

Methods

Continental Chile extends over 38.5 degrees of latitude (17.5°–56°S; 4300 km), and administratively, the country is divided into 16 regions (which range from 15,403 to 132,291 km² in size) and 56 provinces, sequentially ordered from north to south. This arrangement is closely correlated with increasing precipitation and decreasing temperatures with increasing latitude (di Castri and Hajek 1976; Luebert and Plissock 2006). This establishes a smooth gradient in climatic conditions

and a sequence of biomes, from hyperdesert in the north, a Mediterranean climate region in the center and temperate rain forest and cold sub-Antarctic wetlands in the south. In this physical pattern, most of the human population, which is associated with greater environmental alterations, is distributed mainly in the Mediterranean area. In relation to marine systems, the oceanographic conditions of the Chilean coast are strongly influenced by the Humboldt Current System and the Cape Horn Current (Camus 2001; Thiel et al. 2007). To the north of 42°S, the Chilean coast is virtually a line, with few geographical features, but strongly influenced by diverse factors, such as upwelling and El Niño Southern Oscillation (ENSO). On the contrary, south of 42°S is characterized by the high occurrence of geographical accidents, and low salinities due to the influence of fjords (Camus 2001, 2008; Thiel et al. 2007).

We performed an exhaustive bibliographic revision and used herbarium and zoological collections to identify all the species recorded as naturalized in continental Chile and adjacent marine habitats for each of the eight taxonomic groups: a) vascular plants (terrestrial and aquatic); b) nonvascular plants [Bryophyta (mosses), Marchantiophyta (liverworts), and Anthocerotophyta (hornworts)]; c) marine and freshwater macro and micro algae; d) fungi; e) terrestrial vertebrate fauna (amphibians, reptiles, birds and mammals); f) insects; g) marine and freshwater vertebrates (Pisces) and invertebrates (polychaetes and mollusks); and h) terrestrial gastropods. For each taxonomic group, we used the most accepted and comprehensive definition of naturalized species (see Table 1 for details). In general terms, naturalized species were considered as those that are not native to Chile (i.e. nonindigenous) and whose presence is due to intentional or accidental introduction as a result of human activities. We considered all organisms that are naturalized or invasive, but not those that survive only with human assistance (Richardson et al. 2000). The preliminary list of each taxonomic group was then verified by experts (local and international). Thus, we had a second opinion regarding the inclusion or rejection of naturalized species in the database. For each species, we made a substantial effort to compile and organize a database integrating both species characteristics and spatial distribution information (see Table 2 for details). We checked all records for their scientific names and spatial distributions.

To construct the invasion curves of naturalized species we modified the procedure in Pyšek and Prach (1993), following Fuentes et al. (2008). We plotted the cumulative number of species collected/recorded in Chile against time in 20-year periods. Data on the first-year records were gathered from various sources (including online databases, scientific peer-reviewed publications, reports and books) and analyzed for 20-year periods. For invasion curves and the annual rate of first records, we only included species that were first reported in a known year. Thus, terrestrial gastropods, as well as marine and freshwater vertebrates, were excluded from these analyses.

Table I. Definitions of naturalized alien species for each taxonomic group included in the inventory.

Taxonomic group	Definition
Naturalized vascular plants (terrestrial and aquatic)	We included all naturalized alien species <i>sensu</i> Richardson et al. (2000). Additionally, we used expert criteria when the status of the plant species was ambiguous or not supported by publications. Due to a lack of knowledge regarding the native ranges of several species in southern South America (i.e. among Chile, Argentina, Perú, and Bolivia) and to avoid problems in relation to their status (i.e. naturalized or native), we deliberately excluded plants whose natural distribution range fell within the neighboring regions and shared an immediate border with Chile. For this group, the database was constructed on the most recent research by Fuentes et al. (2013), and updated to include new records and spatial information.
Naturalized nonvascular plants (Bryophyta (mosses), Marchantiophyta (liverworts) and Anthocerotophyta (hornworts))	To classify naturalized species in this group we utilized six criteria following Crundwell (1985): 1) absence of subfossil record, 2) evidence of a change in geographical distribution, 3) anomalous geographical distribution, on a world scale or locally, 4) association with some means of introduction e.g. botanic garden or ports, 5) less than the normal amount of genetic variation in populations. Sometimes in dioecious species only one sex is present, and 6) association with open, disturbed or temporary habitats.
Naturalized marine and freshwater macro and microalgae	For both marine and freshwater environment, we included all naturalized species referring to organisms that have established a self-sustaining population, but have not necessarily been involved in an invasion process (Richardson et al. 2000; Falk-Petersen et al. 2006). We considered 12 criteria proposed by different authors to identify species introduced into marine systems worldwide (Chapman and Carlton 1991; Cranfield et al. 1998; Ribera and Boudouresque 1995; Boudouresque and Verlaque 2002), and these were: 1) new report in one area, 2) geographical discontinuity in its worldwide distribution, 3) expansion of its distribution range in the invaded area from its point of introduction following a logical pattern (e.g. gradual decrease in abundance from its point of introduction, colonization of new sites following a spatio-temporal pattern), 4) very localized distribution in the introduced region, restricted when compared with similar native species (for example, of the same genus, of the same functional group, with similar life histories), 5) proximity to the probable source of introduction (e.g. ports, cultivation centers), 6) association (or dependence) with other introduced species, 7) association (or dependence) with anthropic activities (e.g. colonizes docks or other artificial substrates), 8) no native evolutionary origin (e.g. absence of congeners in the variable range), 9) tendency to generate massive proliferations (at least seasonally), 10) is identified as naturalized or invasive in other parts of the world, 11) life stories that infer a high invasive power, and 12) genetically identical distant populations.
Naturalized fungi	For this group, the concept of naturalized species is not easily handled due to the lack of larger inventories of Chilean fungi before 1945. Also, the deficient knowledge in terms of the biogeography of fungi can make it difficult to determine what is a naturalized species. Priority was given to published records of biotrophic and saprotrophic specialists associated to allochthonous plants (trees), apart from collection specimens (CONC-F, Universidad de Concepción) recorded between 2004 and 2017 for the first time in Chile. The preliminary list was revised and completed by external experts.
Naturalized terrestrial vertebrates (mammals, birds, amphibians and reptiles)	In this taxonomic group, we included all naturalized species based on the definition proposed by the United Nations Development Program (UNDP). This definition includes all naturalized alien species that were introduced intentionally or accidentally by humans, establishing a self-sustaining population, without intervention by humans. This general definition was complemented by specific literature for this taxonomic group (e.g. Daniels and Corbett 2003; Lever 1994).
Naturalized insects	For this group, we included all naturalized alien species, referring to organisms that have established a self-sustaining population, but not necessarily ended up in an invasion process (Falk-Petersen et al. 2006; Carvallo 2009).
Naturalized marine and freshwater vertebrates and invertebrates (polychaetes and mollusks) and terrestrial gastropoda	We used the criteria proposed by Orensanz et al. (2002), Castilla et al. (2005) and Castilla and Neill (2009), which states the following: 1) status determined by scientific literature or expert criteria, 2) anomalous geographical distribution, on a world scale or locally, 3) species with wide geographic distribution, including cosmopolitan species, 4) species described as nonindigenous in Chile, and 5) species that are abundant near to ports or aquaculture centers, but rare or not present in other areas of the country.

Table 2. Species traits, descriptions, and traits level included in the inventory of naturalized species present in Chile.

	Plant traits	Description
Taxonomic information	Family, genus, scientific name and author	Text
Distribution	Native range	Name of the countries, continents or bioclimatic zones
	Administrative Regions occupied by the naturalized species in Chile	15= AYP, Arica y Parinacota 1= TAR, Tarapacá 2= ANT, Antofagasta 3= ATA, Atacama 4= COQ, Coquimbo 5= VAL, Valparaíso 13= RME, Región Metropolitana 6= LBO, Libertador Bernardo O'Higgins 7= MAU, Maule 8= BIO, Biobío 9= ARA, La Araucanía 14= LRI, Los Ríos 10= LLA, Los Lagos 11 AIS, Aisén 12= MAG, Magallanes ND = no data.
Introduction	First year report	The year (or range) of the first report in Chile
	Type of introduction	Accidental / Intentional / Other
Impacts	Description of impacts	The known impacts produced by the species around the world.

Results

We identified 1,122 species, which we consider as naturalized at least in some parts of continental Chile and adjacent marine habitats (Table 3). These comprise 790 vascular plants (terrestrial and aquatic); 31 nonvascular plants (Bryophyta, Marchantiophyta and Anthocerotophyta); 18 marine and freshwater macro and micro algae; 71 fungi; 39 terrestrial vertebrates (amphibians, reptiles, birds and mammals); 108 insects; 37 marine and freshwater vertebrates and invertebrates (6 polychaetes, 3 mollusks and 28 Pisces); and 28 terrestrial gastropods (Table 3). For most taxonomic groups, the highest numbers of naturalized species were found mainly in the Mediterranean and rainy climatic zones (Table 3, Fig. 1A, B). Terrestrial vertebrates also showed a significant number of species in the cold steppe zone (Table 3), as did insects, and vascular plants in the semi-arid zone (Table 3, Fig. 1A, B).

The cumulative number of naturalized species collected in Chile over time (i.e. invasion curves) has shown a consistent increment in all taxonomic groups during the 19th and 20th century (Fig. 2). Despite the temporal differences among taxonomic groups in their first records, slopes of invasion curves for naturalized species did differ among taxonomic groups, suggesting different rates of species accumulations. While vascular plants, nonvascular plants, and terrestrial vertebrates showed a slight, but steady increase in the number of naturalized species over the last 150 years (Fig. 2A, B, F), the cumulative number of naturalized species in the remaining four taxa increased steeply at the beginning of the 20th century (Fig. 2C, D, E, G). The total rate of first records remained low between 1850 and 1950 (on average 2.8 first records annually, Fig 3A). Since 1951, first records have increased steeply (on average 5.9 first records annually, Fig 3A). During this period, the continuous increment in first record rates has been consistent in vascular plants, insects and fungi (Fig. 3B). On the contrary, nonvascular plants, terrestrial vertebrates, algae, and marine invertebrates have shown markedly low first record rates over the last 160 years (less than 1 first record annually, Fig. 3B).

Table 3. Number of naturalized species by climatic zones in continental Chile and adjacent marine habitats. The total number of species within each taxonomic group is given in parentheses. Marine organisms are present in the coastal area in front of each climatic zones.

Climatic zones	Vascular plants (790)	Non-vascular plants (31)	Marine & freshwater macro- & microalgae (18)	Fungi (71)	Terrestrial vertebrates (39)	Insects (108)	Marine & freshwater vertebrates & invertebr. (37)	Terrestrial gastropods (28)
Desert	192	0	4	1	13	38	6	1
Semi-arid	318	8	5	1	15	61	10	6
Mediterranean	636	29	11	55	31	108	25	9
Temperate rainy	463	14	13	41	20	78	12	10
Cold steppe	255	8	6	1	23	24	14	2




Figure 1. Total number of naturalized species (**A**) and number of naturalized species per log(area) (**B**) by climatic zones of continental Chile and adjacent marine habitats.

Discussion

Unlike previous studies that have provided a database of naturalized species in Chile (e.g., Fuentes et al. 2013 vascular plants; Moreno et al. 2006 boring polychaetes; Turon et al. 2016 ascidians; Urrutia et al. 2017 aquatic plants, Villaseñor-Parada et al. 2018 marine seaweed and Castilla et al. 2005; Moreno et al. 2006; Castilla and Neill 2009 for marine non-native species), here, we present the first multi-taxa dataset of naturalized species for the whole country. Unfortunately, there are biases in creating this type of inventory because vascular plants are over-represented compared to other taxa, probably because they are more conspicuous, stationary and hence more easily discovered (DAISIE 2009). This plant bias occurred in the present study, where 74.1 % of




Figure 2. Temporal trends in the invasion of naturalized species in continental Chile and adjacent marine habitats, for each taxonomic group for which data on introduction dates were available (terrestrial gastropods are not shown due to lack of data). The cumulative number of recorded species is shown over the last 200 years. **A** vascular plants **B** nonvascular plants **C** marine and freshwater macro and micro algae **D** fungi **E** insects **F** terrestrial vertebrate fauna and **G** marine and freshwater vertebrates and invertebrates.

the naturalized species recorded in Chile are vascular plants. Most of the naturalized species in this database came from herbarium records and zoological collections, while few came from literature. Regarding fauna, there were also some biases in the data sources, e.g. while vertebrates were well represented in museum collections, records of naturalized insects came mainly from literature. The use of natural history collections as the primary source of information highlights their role in naturalized species studies at large spatial and temporal scales in Chile (e.g. see Fuentes et al. 2015 for vascular plants). The present study integrates data in a unique inventory to obtain a more realistic pattern of naturalized species richness and distributions. One major issue came up; when the number of records for certain naturalized species was still particularly low, we could not be completely certain whether these species had naturalized in the area (e.g., nonvascular plants which were poorly sampled before 1950). Nonetheless, the more than 1,100 well-checked species reported in this study should be sufficient to diagnose the situation in Chile in the past and evaluate future risks.

For all taxonomic groups, naturalized species were found to be mainly distributed in Mediterranean and temperate climate regions (central and central-southern Chile), with few species recorded in the far north and south of the country. In the case of vascular plants, several authors have shown similar distribution patterns for naturalized species (Castro et al. 2005; Fuentes et al. 2008; Fuentes et al. 2013). The high concentration of naturalized species in these two climate zones can be related to the fact that since Spanish colonization, most Chileans have lived in this area (ca. 80 % of the population) (Fuentes et al. 2013). Consequently, this area has been strongly trans-




Figure 3. Total temporal trends in first record rates for all species (A) and seven taxonomic groups (B) in Chile. Terrestrial gastropods, as well as marine and freshwater vertebrates are not shown due lack of data.

formed by human activities and land-use changes, while the northern and southern areas have been relatively isolated, mainly due to their climatic conditions and remoteness from the main cities in central-southern Chile (Arroyo et al. 2000; Fuentes et al. 2008, 2015). In marine systems, ship traffic, ballast water and activities associated with aquaculture have been identified as the main introduction vectors for naturalized species (Naylor et al. 2001; Hewitt et al. 2009). In Chile, the main ports are located on the south-central coast (e.g. Valparaíso 33°S, San Antonio 33°35'S, Lirquén 36°42'S, Talcahuano 36°43'S, San Vicente 36°44'S, Coronel 37°01'S and Corral 39°52'S), which increases the susceptibility of these ecosystems to be invaded by naturalized species. Paradoxically, this sector of the Chilean coast has been one of the least studied in terms of biological invasions (Villaseñor-Parada et al. 2017), and even though the number of reports of naturalized species was found to be high, this number could be underestimated due to the lack of information available (Camus 2005; Carlton 2009; Castilla and Neill 2009; Villaseñor-Parada et al. 2017). The few naturalized species recorded in the extreme north and south of the country, could be associated with sampling efforts in these areas, the fact that naturalized species have not yet arrived to these areas due to low propagule pressure or introduction effort, and/or the fact that some species may have arrived, but not thrived because such habitats were incompatible with their ecological niches (Garrido 1985; Jaksic 1998; Fuentes et al. 2015; Villaseñor-Parada et al. 2017, 2018). For vascular plants in particular, the low species density in the Atacama Desert may be attributable to the harsh climatic conditions there, while the lack of naturalized plants in the far south may be the result of a combination of inaccessibility and lower levels of anthropization (Fuentes et al. 2008, 2015). In the case of insects, the highest presence of naturalized species was found to be concentrated in

Chile's central zone, which could be explained by the higher concentration of seaports, border crossings and airports, and the import of biological control agents (Parra and González 2007), pollinators (Montalva et al. 2008), and products of plant or animal origin that, in turn, are contaminated with foreign insects (Estay 2016). Thus, a more comprehensive inventory of naturalized species in these areas will help to identify priority responses to the growing threats from biological invasions.

Thorough documentation of the accumulation of naturalized species, allows us to assess the accumulation process and the dynamics of the establishment and expansion of naturalized species in Chile. We identified invasion periods in Chile based on bibliographic revisions, herbarium and zoological collections for each of the eight taxonomic groups analyzed. Comparatively, the invasion curves show that naturalized species first underwent a positive increment, followed by an apparent plateau phase, mainly in vascular plants, nonvascular plants, and terrestrial vertebrates. In the remaining four taxonomic groups (fungi, insects, marine and freshwater vertebrates and invertebrates, macro and microalgae), the cumulative number of naturalized species increased abruptly at the beginning of the 20th century. The lack of collections before 1900 is also evident for most of the taxonomic groups. In marine systems, the first reports of marine flora and fauna date back to the end of the 18th century, as a result of scientific expeditions that were limited to enumerating the different taxa collected, with few and brief descriptions (Etcheverry 1958; Ramírez 2010). In 1940, the first publications by resident Chilean taxonomists who ratified and added information on the presence of the aforementioned taxa started to appear (Ramírez 2010). Therefore, although the results observed when analyzing the temporal trend of marine invasions show a growing increase since 1900 (Fig. 2C, G), these results are biased due to a lack of prior information before this date. Since the Spanish colonization, an important number of species has been introduced in the drier central Mediterranean climate area, associated with landscape transformations (Aronson et al. 1998). Additionally, in the 19th century there was an important increase in the number of biodiversity records in Chile associated with the inventories developed by naturalists, such as Claudio Gay (Gay 1845, 1854). For vascular plants, there were sustained increments in the collection of weeds from 1894 to 1934, associated with wheat imports (Matthei 1995), whereas the increase of naturalized fungal species, especially macromycetes, is strongly correlated with the fast and extensive spread of naturalized timber plantations, mainly pine and eucalyptus, during the late decades of the 20th century (Palfner and Casanova 2019). The latter authors observed an almost threefold increase in allochthonous, and ubiquitous fungi associated with *Pinus radiata* plantations in central-southern Chile within the last 40 years. For other taxonomic groups, such as terrestrial vertebrates, the record of introduced species is relatively more complete and the relative lower rate of increase may be due to control measures implemented by government services (Iriarte et al. 2005). Specific phytosanitary regulations and other laws related to the introduction of naturalized species have most likely worked in these cases.

Considering naturalized species as a whole, this inventory highlights the consistent increase in the rate of new naturalizations since 1950, especially for some taxonomic groups, such as insects, fungi, and vascular plants. Even where rates of establishment

did not prove to be rising for nonvascular plants, algae, marine, and terrestrial vertebrates, the cumulative number of naturalized taxa did show a consistent increase for these taxonomic groups. Our results are consistent with the fact that the annual rate of first records worldwide has increased during the last 200 years, with 37% of all first records reported recently (1970–2014) (Seebens et al. 2017). Continuous increases in the rates of naturalized species first records in Chile, stress the need to improve the implementation of national legislation and international agreements that aim to reduce the threats that naturalized alien species pose to biodiversity, particularly in the case of insects and fungi. Additionally, the above pattern suggests that the number of new naturalized species will most likely continue to increase because current tools to prevent biological invasions are not effective enough to slow down the increment in the number of naturalized species (Seebens et al. 2017). This inventory is a basis for future studies that analyze more detailed biological patterns and mechanisms that explain invasion processes in many taxa that have been understudied (*sensu*, Capdevila-Argüelles and Zilletti 2005). In this context, to effectively assess consequences and risks, in terms of the spread and invasion of individual species, documentation with an efficient combination of species records requires a coordinated effort across multiple government agencies and research institutions, in addition to well-designed and specifically oriented sampling and monitoring programs.

Acknowledgements

A previous version of the current database was created during the study conducted by the University of Concepción under the contract SCT N°20/2015 “Consultoría para elaborar el catálogo de las especies exóticas asilvestradas/naturalizadas en Chile, en el marco del Proyecto GEF/MMA/PNUD EEI AJF”. We would like to thank the Ministerio del Medio Ambiente de Chile (MMA) and PNUD, especially Fernando Baeriswyl, Charif Tala, Emma Elgueta, Macarena Isla and Giuliana Furci. We are grateful to Fernando Carrasco, Héctor Ibarra (terrestrial vertebrates) and Fabio Labra (Facultad de Ciencias, Universidad San Tomás, Chile) for their help during the Project. NF and AS are supported by FONDECYT N°1181688, AP and PS funded by CONICYT PIA grant AFB 170008. We thank all the people at CONC Herbarium (University of Concepción).

References

- Armesto JJ, Rozzi R, Smith-Ramírez C, Arroyo MTK (1998) Conservation targets in South American temperate forests. *Science* 5392: 1271–1272. <https://doi.org/10.1126/science.282.5392.1271>
- Aronson J, Del Pozo A, Ovalle C, Avendano J, Lavin A, Etienne M (1998) Land use changes and conflicts in Central Chile. In: Rundel PW, Montenegro G, Jaksic F (Eds) *Landscape disturbance and biodiversity in Mediterranean type ecosystems*. Springer-Verlag Germany, 155–168. https://doi.org/10.1007/978-3-662-03543-6_9

- Arroyo MTK, Marticorena C, Matthei O, Cavieres L (2000) Plant invasions in Chile: present patterns and future predictions. In: Mooney A, Hobbs R (Eds) *Invasive species in a changing world*. Island Press, California USA, 385–421.
- Boudouresque CF, Verlaque M (2002) Biological pollution in the Mediterranean Sea: invasive versus introduced macrophytes. *Marine Pollution Bulletin* 44: 32–38. [https://doi.org/10.1016/S0025-326X\(01\)00150-3](https://doi.org/10.1016/S0025-326X(01)00150-3)
- Camus PA (2001) Biogeografía marina de Chile continental. *Revista Chilena de Historia Natural* 74: 587–617. <https://doi.org/10.4067/S0716-078X2001000300008>
- Camus PA (2005) Introducción de especies en ambientes marinos chilenos: no solo exóticas, no siempre evidentes. *Revista Chilena de Historia Natural* 78: 155–159. <https://doi.org/10.4067/S0716-078X2005000100011>
- Camus PA (2008) Diversidad, distribución y abundancia de especies en ensamblajes intermareales rocosos. *Revista de Biología Marina y Oceanografía* 43: 615–627. <https://doi.org/10.4067/S0718-19572008000300021>
- Capdevila-Argüelles L, Zilletti B (2005) *Issues in Bioinvasion Science*. Springer, Netherlands.
- Carlton JT (2009) Deep invasion ecology and the assembly of communities in historical time. In: Rilov G, Crooks JA (Eds) *Biological Invasions in Marine Ecosystems: Ecological, Management, and Geographic Perspectives*. Springer-Verlag, Berlin, 13–56. https://doi.org/10.1007/978-3-540-79236-9_2
- Carvalho GO (2009) Especies exóticas e invasiones biológicas. *Ciencia Ahora* 23(12): 15–21.
- Castilla JC, Uribe M, Bahamonde N, Clarke M, Desqueyroux-Faúndez R, Kong I, Moyano H, Rozbaczylo N, Santelices B, Valdovinos C, Zavala P (2005) Down under the southeastern Pacific: marine non-indigenous species in Chile. *Biological Invasions* 7: 213–232. <https://doi.org/10.1007/s10530-004-0198-5>
- Castilla JC, PE Neill (2009) Marine Bioinvasions in the Southeastern Pacific: Status, Ecology, Economic Impacts, Conservation and Management. In: Rilov G, Crooks JA (Eds) *Biological Invasions in Marine Ecosystems: Ecological, Management, and Geographic Perspectives*. Springer-Verlag, Berlin, 439–457. https://doi.org/10.1007/978-3-540-79236-9_26
- Castro SA, Figueroa J, Muñoz-Schick M, Jaksic FM (2005) Minimum residence time, biogeographical origin, and life cycle as determinants of the geographical extent of naturalized plants in continental Chile. *Diversity and Distributions* 11: 183–191. <https://doi.org/10.1111/j.1366-9516.2005.00145.x>
- Chapman JW, Carlton JT (1991) A test of criteria for introduced species: the global invasion by the isopod *Synidotea laevidorsalis* (Miers, 1881). *Journal of Crustacean Biology* 11: 386–400. <https://doi.org/10.2307/1548465>
- Cranfield HJ, Gordon DP, Willan RC, Marshall BA, Battershill CN, Francis MP, Nelson WA, Glasby CJ, Read GB (1998) Adventive marine species in New Zealand. Wellington. <http://docs.niwa.co.nz/library/public/NIWATR34.pdf>
- Crundwell AC (1985) The introduced bryophytes of the British Isles. *Bulletin of the British Bryological Society* 45: 8–9.
- DAISIE (Ed.) (2009) *Handbook of alien species in Europe*. Springer, Berlin.
- Daniels MJ, Corbett L (2003) Redefining introgressed protected mammals: when is a wildcat a wild cat and a dingo a wild dog? *Wildlife Research* 30: 213–218. <https://doi.org/10.1071/WR02045>

- di Castri F, Hajek ER (1976) *Bioclimatología de Chile*. Santiago Chile. Vicerrectoría Académica de la Universidad Católica de Chile. https://www.ecolya.cl/documentos/bioclimatologia_de_chile.pdf
- Estay S (2016) Invasive insects in the Mediterranean forests of Chile. In: Paine T, Lieutier F (Eds) *Insects and diseases of Mediterranean forest systems*. Springer, Cham, 379–396. https://doi.org/10.1007/978-3-319-24744-1_13
- Etcheverry H (1958) Bibliografía de las algas chilenas. *Revista de Biología Marina* 7: 163–182.
- Falk-Petersen J, Bøhn T, Sandlund OT (2006) On the numerous concepts in invasion biology. *Biological Invasions* 8: 1409–1424. <https://doi.org/10.1007/s10530-005-0710-6>
- Fuentes N, Ugarte E, Kühn I, Klotz S (2008) Alien plants in Chile: inferring invasion periods from herbarium records. *Biological Invasions* 10: 649–657. <https://doi.org/10.1007/s10530-007-9159-0>
- Fuentes N, Ugarte E, Kühn I, Klotz S (2010) Alien plants in southern South America. A framework for evaluation and management of mutual risk of invasion between Chile and Argentina. *Biological Invasions* 12: 3227–3236. <https://doi.org/10.1007/s10530-010-9716-9>
- Fuentes N, Pauchard A, Sanchez P, Esquivel J, Marticorena A (2013) A new comprehensive database of alien plant species in Chile based on herbarium records. *Biological Invasions* 15: 847–858. <https://doi.org/10.1007/s10530-012-0334-6>
- Fuentes N, Saldaña A, Kühn I, Klotz S (2015) Climatic and socio-economic factors determine the level of invasion in Chile. *Plant Ecology and Diversity* 8: 371–377. <https://doi.org/10.1080/17550874.2014.984003>
- Gardener M, Bustamante RO, Herrera I, Durigand G, Pivello VR, Moro MF, Stoll A, Langdon B, Baruch Z, Rico A, Arredondo-Núñez A, Flores S (2012) Plant invasion in Latin America: fast track to a more focused agenda. *Plant Ecology and Diversity* 5: 225–232. <https://doi.org/10.1080/17550874.2011.604800>
- Garrido N (1985) *Index Agaricalium Chilensium*. Bibliotheca Mycologica 99. Vaduz J Cramer.
- Gay C (1845) *Historia Física y Política de Chile*. Tomo I. Thunot Paris, France.
- Gay C (1854) *Atlas de la Historia Física y Política de Chile*. Tomo II. Thunot Paris, France.
- Habit E, Dyer B, Vila I (2006) Estado de conocimiento de los peces dulceacuícolas de Chile. *Gayana* 70(1):100–113. <https://doi.org/10.4067/S0717-65382006000100016>
- Hewitt CL, Gollasch S, Minchin D (2009) The vessel as a vector - biofouling, ballast water and sediments. In: Rilov G, Crooks JA (Eds) *Biological Invasions in Marine Ecosystems: Ecological, Management, and Geographic Perspectives*. Ecological Studies Series. Springer-Verlag, 117–131. https://doi.org/10.1007/978-3-540-79236-9_6
- Hulme PE (2009) Trade, transport and trouble: managing invasive species pathways in an area of globalization. *Journal of Applied Ecology* 46: 10–18. <https://doi.org/10.1111/j.1365-2664.2008.01600.x>
- Iriarte JA, Lobos G, Jaksic FM (2005) Invasive vertebrate species in Chile and their control and monitoring by governmental agencies. *Revista Chilena de Historia Natural* 78: 149–154. <https://doi.org/10.4067/S0716-078X2005000100010>
- Jaksic FM (1998) Vertebrate invaders and their ecological impacts in Chile. *Biodiversity & Conservation* 7(11): 1427–1445. <https://doi.org/10.1023/A:1008825802448>
- Lever C (1994) *Naturalized Animals: The Ecology of Successfully Introduced Species*. Poyser Natural History, London.

- Luebert F, Pliscoff P (2006) Sinopsis bioclimática y vegetal de Chile. Santiago de Chile. Editorial Universitaria.
- Mack RN, Simberloff D, Lonsdale WM, Evans H, Clout M, Bazzaz FA (2000) Biotic invasions: causes, epidemiology, global consequences, and control. *Ecological Applications* 10(3): 689–710. [https://doi.org/10.1890/1051-0761\(2000\)010\[0689:BICEGC\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2000)010[0689:BICEGC]2.0.CO;2)
- Matthei O (1995) Manual de las malezas que crecen en Chile. Concepción (Chile). Alfabeto Impresores.
- Mihulka S, Pyšek P (2001) Invasion history of *Oenothera* congeners in Europe: a comparative study of spreading rates in the last 200 years. *Journal of Biogeography* 28:597–609. <https://doi.org/10.1046/j.1365-2699.2001.00574.x>
- Mittermeier RA, Robles Gil P, Hoffman M, Pilgrim J, Brooks T, Mittermeier CG, Lamoreux J, Da Fonseca GAB (2005) Hotspots revisited: earth's biologically richest and most endangered terrestrial ecoregions. University of Chicago Press for Conservation International. USA.
- Montalva J, Arroyo MK, Ruz L (2008) *Bombus terrestris* Linnaeus (Hymenoptera: Apidae: Bombini) en Chile: Causas y consecuencias de su introducción. *Revista del Jardín Botánico Chagual* Año 6: 6–5.
- Moreno RA, Neill PE, Rozbaczylo N (2006) Native and non-indigenous boring polychaetes in Chile: a threat to native and commercial mollusc species. *Revista Chilena de Historia Natural* 79: 263–278. <https://doi.org/10.4067/S0716-078X2006000200012>
- Myers N, Mittermeier RA, Mittermeier CG, Da Fonseca GAB, Kent J (2000) Biodiversity hotspots for conservation priorities. *Nature* 403: 853–858. <https://doi.org/10.1038/35002501>
- Naylor RL, Williams SL, Strong DR (2001) Aquaculture A Gateway for Exotic Species *Science* 294 (5547): 1655–1656. <https://doi.org/10.1126/science.1064875>
- Núñez M, Pauchard A (2010) Biological invasions in developing and developed countries: does one model fit all? *Biological Invasions* 12:707–714. <https://doi.org/10.1007/s10530-009-9517-1>
- Orensanz JM, Schwindt E, Pastorino G, Bortolus A, Casas G, Darrigran G, Elías R, López Gappa JJ, Obenar S, Pascual M, Penchaszadeh P, Piriz ML, Scarabino F, Spivak ED, Valdarino EA (2002) No longer the pristine confines of the world ocean: a survey of exotic marine species in the southwestern Atlantic. *Biological Invasions* 4: 115–143. <https://doi.org/10.1023/A:1020596916153>
- Ormazabal C (1993) The conservation of biodiversity in Chile. *Revista Chilena de Historia Natural* 66:383–402. http://rchn.biologiachile.cl/pdfs/1993/4/Ormazabal_1993.pdf
- Palfner G, Casanova-Katny A (2019) Micocenosis en remanentes de bosque nativo y en plantaciones forestales en la península de Arauco, Biobío, Chile: composición, aspectos funcionales y conservación (Comparison of the Mycobiota in remnants of native forests and forest plantations in the Arauco península of the Biobío region, highlighting functional and conservation aspects). In: Smith-Ramírez C, Squeo FA (Eds) *Biodiversidad y Ecología de los Bosques Costeros de Chile*, Editorial Universidad de Los Lagos: 175–210.
- Parra P, González M (2007) *Megarhyssa praezellens*, Parasitoide de la avispa de la madera *Tremex fuscicornis*, en Chile. *Instituto Forestal de Chile INFOR* 13(3): 473–489.
- Pauchard A, Cavieres L, Bustamante R, Becerra P, Rapoport E (2004) Increasing the understanding of plant invasions in southern South America: first symposium on Alien

- Plant Invasions in Chile. *Biological Invasions* 6:255–257. <https://doi.org/10.1023/B:BINV.0000022137.61633.09>
- Pimentel D, Zuniga R, Morrison D (2005) Update on the environmental and economic costs associated with alien-invasive species in the United States. *Ecological Economics* 52(3): 273–28. <https://doi.org/10.1016/j.ecolecon.2004.10.002>
- PNUD (2017) Catálogo de las especies exóticas asilvestradas/naturalizadas en Chile. Laboratorio de Invasiones Biológicas (LIB) Universidad de Concepción, Proyecto GEF/MMA/PNUD Fortalecimiento de los Marcos Nacionales para la Gobernabilidad de las Especies Exóticas Invasoras: Proyecto Piloto en el Archipiélago de Juan Fernández. Santiago de Chile.
- Pyšek P, Prach K (1993) Plant invasion and the role of riparian habitats: a comparison of four species alien to central Europe. *Journal of Biogeography* 20: 413–420. <https://doi.org/10.2307/2845589>
- Pyšek P, Richardson DM, Pergl J, Jarošík V, Sixtová Z, Weber E (2008) Geographical and taxonomic biases in invasion ecology. *Trends in Ecology and Evolution* 23: 237–244. <https://doi.org/10.1016/j.tree.2008.02.002>
- Pyšek P, Richardson DM (2010) Invasive species, environmental change and management, and health. *Annual Review of Environment and Resources* 35: 25–55. <https://doi.org/10.1146/annurev-environ-033009-095548>
- Quiroz CL, Pauchard A, Cavieres LA, Anderson CB (2009) Análisis cuantitativo de la investigación en invasiones biológicas en Chile: tendencias y desafíos. *Revista Chilena de Historia Natural* 82: 497–505. <https://doi.org/10.4067/S0716-078X2009000400005>
- Ramírez ME (2010) Algas Marinas Bentónicas: Chlorophyta, Ochrophyta (Phaeophyceae) y Rhodophyta. In: Palma S, Báez P, Pequeño G (Eds) *Bibliografía Sobre Biodiversidad Acuática de Chile*. Comité Oceanográfico Nacional, Valparaíso, 13–28. http://www.cona.cl/pub/libro_bibliografia/1.PDF
- Ribera MA, Boudouresque CF (1995) Introduced marine plants, with special reference to macroalgae: mechanisms and impact. In: Round FE, Chapman DJ (Eds) *Progress in Phycological Research*. Biopress Bristol, 187–268. https://people.mio.osupytheas.fr/~boudouresque/Publications_pdf/Ribera_Boudouresque_1995_Prog_Phycol_Res.pdf
- Richardson DM, Pyšek P, Rejmánek M, Barbour MG, Panetta FD, West CJ (2000) Naturalization and invasion of alien plants: concepts and definitions. *Diversity and Distributions* 6: 93–107. <https://doi.org/10.1046/j.1472-4642.2000.00083.x>
- Rodríguez R, Marticorena C, Alarcón D, Baeza C, Cavieres L, Finot VL, Fuentes N, Kiessling A, Mihoc M, Pauchard A, Ruiz E, Sanchez P, Marticorena A (2018) Catálogo de las plantas vasculares de Chile. *Gayana Botánica* 75(1): 1–430. <https://doi.org/10.4067/S0717-66432018000100001>
- Seebens H, Blackburn TM, Dyer EE, Genovesi P, Hulme PE, Jeschke JM, Pagad S, Pyšek P, Winter M, Arianoutsou M, Bacher S, Blasius B, Brundu G, Capinha C, Celesti-Grapow L, Dawson W, Dullinger S, Fuentes N, Jäger H, Kartesz J, Kenis M, Kreft H, Kühn I, Lenzner B, Liebhold A, Mosena A, Moser D, Nishino M, Pearman D, Pergl J, Rabitsch W, Rojas-Sandoval J, Roques A, Rorke S, Rossinelli S, Roy HE, Scalera R, Schindler S, Štajerová K, Tokarska-Guzik B, van Kleunen M, Walker K, Weigelt P, Yamanaka T, Essl F (2017) No

- saturation in the accumulation of alien species worldwide. *Nature Communications* 8: 14435. <https://doi.org/10.1038/ncomms14435>
- Speziale K, Lambertucci S, Carrete M, Tella J (2012) Dealing with non-native species: what makes the difference in South America. *Biological Invasions* 14: 1609–1621. <https://doi.org/10.1007/s10530-011-0162-0>
- Thiel M, Macaya EC, Acuña E, Arntz WE, Bastias H, Brokordt K, Camus PA, Castilla JC, Castro LR, Cortés M, Dumont CP, Escribano R, Fernández M, Gajardo JA, Gaymer CF, Gomez I, González AE, González HE, Haye PA, Illanes JE, Iriarte JL, Lancelloti DA, Luna-Jorquera G, Luxoro C, Manriquez PH, Marín V, Muñoz P, Navarretes SA, Perez E, Poulin E, Sellanes J, Sepúlveda HH, Stotz W, Tala F, Thomas A, Vargas CA, Vasquez JA, Vega JMA (2007) The Humboldt current system of northern and central Chile Oceanographic processes, ecological interactions and socioeconomic feedback. *Oceanography and Marine Biology: An Annual Review* 45: 195–344. <https://doi.org/10.1201/9781420050943.ch6>
- Turon X, Cañete JI, Sellanes J, Rocha RM, López-legentil S (2016) Too cold for invasions? Contrasting patterns of native and introduced ascidians in subantarctic and temperate Chile. *Management of Biological Invasions* 7: 77–86. <https://doi.org/10.3391/mbi.2016.7.1.10>
- Urrutia J, Sánchez P, Pauchard A, Hauenstein E (2017) Plantas acuáticas invasoras presentes en Chile: Distribución, rasgos de vida y potencial invasor. *Gayana Botánica* 74(1): 147–157. <https://doi.org/10.4067/S0717-66432017005000324>
- van Kleunen M, Dawson W, Essl F, Pergl J, Winter M, Weber E, Kreft H, Weigelt P, Kartesz J, Nishino M, Antonova LA, Barcelona JF, Cabezas FJ, Cárdenas D, Cárdenas-Toro J, Castaño N, Chacón E, Chatelain C, Ebel AL, Figueiredo E, Fuentes N, Groom QJ, Henderson L, Inderjit, Kupriyanov A, Masciadri S, Meerman J, Morozova O, Moser D, Nickrent DL, Patzelt A, Pelter PB, Baptiste MP, Poopath M, Schulze M, Seebens H, Shu W-S, Thomas J, Velayos M, Wieringa JJ, Pyšek P (2015) Global exchange and accumulation of non-native plants. *Nature* 525: 100–103. <https://doi.org/10.1038/nature14910>
- Vidal MA, Díaz-Páez H (2012) Biogeography of Chilean herpetofauna: biodiversity hotspot and extinction risk. In: Stevens L (Ed.) *Global advances in biogeography*. Intech Editions, Flagstaff, USA. http://cdn.intechopen.com/pdfs/34659/InTech-Biogeography_of_chilean_herpetofauna_biodiversity_hotspot_and_extinction_risk.pdf
- Villaseñor-Parada C, Pauchard A, Macaya EC (2017) Ecología de invasiones marinas en Chile continental: ¿Qué sabemos y que nos falta por saber? *Revista de Biología Marina y Oceanografía*, 52(1): 01–17. <https://doi.org/10.4067/S0718-19572017000100001>
- Villaseñor-Parada C, Pauchard A, Ramírez M, Macaya EC (2018) Macroalgas exóticas en la costa de Chile: patrones espaciales y temporales en el proceso de invasión. *Latin American journal of aquatic research* 46(1): 147–165. <https://doi.org/10.3856/vol46-issue1-fulltext-15>

Supplementary material I

A comprehensive inventory of naturalized species in Chile

Authors: Nicol Fuentes, Alicia Marticorena, Alfredo Saldaña, Viviane Jerez, Juan Carlos Ortiz, Pedro Victoriano, Rodrigo A. Moreno, Juan Larraín, Cristobal Villaseñor-Parada, Götz Palfner, Paulina Sánchez, Aníbal Pauchard

Data type: dataset

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/neobiota.60.55366.suppl1>